

EXPECTORANT SYRUP



taking it.

Side effects:

Common adverse effects are as follows:
Constipation; diarrhea; dizziness; drowsiness; excitability; headache; loss of appetite; nausea; nervousness or anxiety; trouble sleeping; upset stomach; vomiting; weakness

Contraindications:

history of adrenal gland problems (eg, adrenal gland tumor); heart problems; high blood pressure; diabetes; blood vessel problems; stroke; glaucoma; a blockage of your bladder, stomach, or intestines; ulcers; trouble urinating; an enlarged prostate or other prostate problems; seizures; or an overactive thyroid

Interactions:

- Beta-blockers (eg, propranolol), catechol-O-methyltransferase (COMT) inhibitors (eg, tolcapone), furazolidone, indomethacin, MAO inhibitors (eg, phenelzine), sodium oxybate (GHB), or tricyclic antidepressants (eg, amitriptyline) because the side effects of chlorpheniramine/guaifenesin/pseudoephedrine may be increased
- Digoxin or droxidopa because the risk of irregular heartbeat or heart attack may be increased
- Bromocriptine or hydantoins (eg, phenytoin) because side effects may be increased by chlorpheniramine/guaifenesin/pseudoephedrine
- Guanadrel, guanethidine, metcamylamine, methyldopa, or reserpine because their effectiveness may be decreased by chlorpheniramine/guaifenesin/pseudoephedrine

Administration:

- Chlorpheniramine/guaifenesin/pseudoephedrine may be taken with or without food.
- If you miss a dose of this drug, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once.
- Chlorpheniramine/guaifenesin/pseudoephedrine may cause dizziness, drowsiness, or blurred vision. Do not drive, operate machinery, or do anything else that could be dangerous.
- Using chlorpheniramine/guaifenesin/pseudoephedrine alone, with certain other medicines, or with alcohol may lessen your ability to drive or perform other potentially dangerous tasks.

Dosage Form:


Syrup

Each 5 ml contains 100 mg Guaifenesin + 2 mg Chlorpheniramine + 30 mg Pseudoephedrine

Packaging: 60 ml bottle

References:

- 1) Lexicomp, s Drug Reference Handbooks, American Pharmacists Association, 20th edition, page 347
- 2) <https://www.drugs.com/cdi/chlorpheniramine-guaifenesin-pseudoephedrine.html>
- 3) <http://apps.who.int/medicinedocs/documents/s17501en/s17501en.pdf>
- 4) <http://www.medicinenet.com/guaifenesindecongestantantihistamine-oral/article.htm>
- 5) <http://www.healthguidance.org/entry/17015/1/Side-Effects-of-Expectorants.html>
- 6) <http://reference.medscape.com/drug/sudafed-nexafed-pseudoephedrine-343412>
- 7) <http://reference.medscape.com/drug/chlortrimeton-chlorpheniramine-343386>
- 8) <http://reference.medscape.com/drug/mucinex-organidin-nr-guaifenesin-343403>



Indications:

Relieving symptoms of sinus congestion, runny nose, sneezing, and throat and airway irritation due to colds, upper respiratory infections, and allergies.

Pharmacological Category:

Expectorant-antihistamine-decongestant

Pregnancy:

If you become pregnant while taking this drug, discuss with your doctor the benefits and risks of using.

Lactation:

It is unknown if this drug is excreted in breast milk. Do not breast-feed while