Lito Kharazmi

Salbutamol

Salbutamol Syrup

Dosage & Indications:
Pharmacologic category:
Bronchodilator

Treatment bronchospasm:
Children 2–6 years: 0.25–0.5 ml, every 6–8 hours. Maximum dose: 4 mg daily.
Children 6–14 years: 5 ml, every 6–8 hours. Maximum dose 24 mg daily.
Children>14 years and adults: 2–4 mg, every 6–8 hours. Maximum dose: 32 mg, daily.

Contraindications:
Hypersensitivity to salbutamol or any of the other ingredients.

Pregnancy & Breast-feeding:
Use with caution in pregnancy. Inhaled drugs for asthma can be taken as normal during breast-feeding.

Warnings:
Use with caution in children under 2 years of age.

Cautions:
Salbutamol should use with caution in patients with cardiac disease. Administer with meals to minimize gastric irritation

Interactions:
Amitriptyline, clomipramine, desipramine, doxepin, imipramine, isocarboxazid, linezolid, maprotiline, nortriptyline, phenelzine, tranylcypromine, trazodone.

Side effects:
Lactic acidosis (with high doses), nausea, tremor, nervousness in children aged 2–6 years, fever, vomiting, headache, dizziness, cough.

Date of last authorization and revision:
February 2019

ATW.065/LF.01/E

لیتو خوارزمی

سالبوتامول

شربت سالبوتامول

موارد و مقدار مصرف:
دسته دارویی:گشاد کننده برونش

درمان برونکواسپاسم:
کودکان ۶-۲ سال: ۰/۵-۰/۲۵ میلی لیتر هر ۸-۶ ساعت.حداکثر دوز مجاز روزانه: ۴ میلی گرم.
کودکان ۱۴-۶ سال: ۵ میلی لیتر هر ۸-۶ ساعت. حداکثر دوز مجاز روزانه: ۲۴ میلی گرم.
کودکان بالای ۱۴ سال و بزرگسالان: ۴-۲ میلی گرم هر ۸-۶ ساعت، حداکثر دوز مجاز روزانه: ۳۲ میلی گرم.

موارد منع مصرف:
حساسیت به سالبوتامول یا هر یک از اجزای دارو.

بارداری و شیردهی:
در دوران بارداری با احتیاط مصرف شود.
در دوران شیردهی بهتر است از فرم استنشاقی سالبوتامول استفاده شود.

هشدارها:
درکودکان زیر ۲ سال با احتیاط مصرف شود.

احتیاط های مصرف:
در بیماران قلبی با احتیاط مصرف شود.
برای کاهش عوارض گوارشی، بهتر است دارو همراه با غذا مصرف شود.

تداخل های دارویی:
آمی تریپتیلین، کلومیپرامین، دزیپرامین، دوکسپین، ایمیپرامین، ایزوکربوکسازید، لینزولید، ماپروتیلین، نورتریپتیلین، فنلزین، ترانیل سیپرومین، ترازودون.

عوارض جانبی:
اسیدوز لاکتیک (با دوزهای بالا)، تهوع، لرزش بدن، تحریک عصبی در کودکان ۶-۲ سال، تب، استفراغ، سردرد، سرگیجه، سرفه.

تاریخ آخرین تدوین و بازبینی:
اسفند ۱۳۹۷
صدای مشتری: ۰۲۱-۴۴۵۴۵۴۲۰

ATW.065/LF.01/P